

Braid group actions and tensor products for Yangians

Yilan Tan

Abstract

We introduce a braid group action on l -tuple of rational functions for the finite-dimensional representations of Yangians $Y(\mathfrak{g})$, where $\mathfrak{g}$ is a complex simple Lie algebra. It provides an efficient way to compute certain polynomials which allows us to provide a finite set of numbers at which the tensor product of Kirillov-Reshetikhin modules of Yangians may fail to be cyclic.

Key words: Yangians; braid group action; tensor product; cyclic condition.

1 Introduction

Let $\mathfrak{g}$ be a complex simple Lie algebra of rank l . For each $\mathfrak{g}$, there exists a Hopf algebra $Y(\mathfrak{g})$, called the Yangian of $\mathfrak{g}$, which is a deformation of the universal enveloping algebra of the current algebra $\mathfrak{g}[t]$. For each l -tuple of polynomials $\pi = (\pi_1(u), \dots, \pi_l(u))$, Drinfeld showed in [9] that there exists a unique finite-dimensional irreducible representation $V(\pi)$ of the Yangian $Y(\mathfrak{g})$. If there exists an $i \in I = \{1, \dots, l\}$ such that $\pi_i(u) = (u - a)(u - (a + 1)) \dots (u - (a + m - 1))$ and $\pi_j(u) = 1$ for $j \neq i$, then $V(\pi)$ is called the Kirillov-Reshetikhin module. In this case, π is denoted by $\pi_{a,m}^{(i)}$. In particular, when $m = 1$, $V(\pi)$ is called an i -th fundamental representation, denoted by $V_a(\omega_i)$, and the l -tuple of polynomials is denoted by $\pi_{i,a}$. When $\mathfrak{g} = \mathfrak{sl}_2$, $V(\pi)$ can be realized as a tensor product of Kirillov-Reshetikhin modules [6]. For $\mathfrak{g} \neq \mathfrak{sl}_2$, the structure of $V(\pi)$ is less clear. Chari and Pressley showed in [5] that $V(\pi)$ is a sub-quotient of a tensor product of fundamental representations $V_{a_1}(\omega_{b_1}) \otimes V_{a_2}(\omega_{b_2}) \otimes \dots \otimes V_{a_k}(\omega_{b_k})$, where the tensor factors may take any order. In the context of quantum affine algebras $U_q(\hat{\mathfrak{g}})$, the irreducibility of the tensor product of fundamental representations has been studied by many authors who used a variety of methods [1, 10, 12, 14, 19]. This question was generalized by Chari [4], who gave a concrete irreducibility condition for a tensor product of Kirillov-Reshetikhin modules of quantum affine algebras. We note that determining certain polynomials is the key to giving an irreducibility condition for the aforementioned tensor product and Chari connected it with defining a braid group action on the set of l -tuple of rational functions.

Adopting the methods in the Chari's paper, we provided an analogous concrete irreducibility condition for the tensor product of fundamental representations of $Y(\mathfrak{g})$ [16, 17], where $\mathfrak{g}$ is either of classical type or of type G_2 . Since there is no analogous braid group action available for Yangians at that time, we computed certain polynomials using some of the defining relations of $Y(\mathfrak{g})$. In practice, it is tedious to compute certain polynomials in the same way when $\mathfrak{g}$ is an exceptional simple Lie algebra of type either E or F , which is why we had not obtained an irreducibility condition in these cases at that moment. This motivated us to introduce a braid group action for Yangians.

We would like to mention here that it has been established by Varagnolo [18] that together with results of Nakajima [15], $Y(\mathfrak{g})$ and $U_q(\hat{\mathfrak{g}})$ has the same finite-dimensional representation theory. The applications of quiver varieties is essential in these papers. More recently, Gautam and Valerio Toledano [11] provided an equivalence of meromorphic braided tensor categories, where the Yangians and quantum affine algebras endowed with the deformed Drinfeld coproducts. The aim of this paper is to give an independent proof of a cyclicity condition for a tensor product (the

standard coproduct) of Kirillov-Reshetikhin modules of Yangians.

This paper is organized as follows. In Section 2, the Yangian $Y(\mathfrak{g})$ and its finite-dimensional representations are summarized. In Section 3, we define a braid group action on the set of l -tuple of rational functions over complex numbers and show that the action is well defined. In Section 4, we show in Proposition 4.5 that the eigenvalues of $h_i(u)$ on an extremal vector is governed by the braid group action defined in Section 3. Using this, one can find a cyclicity condition for the tensor product of finite-dimensional irreducible representations of $Y(\mathfrak{g})$. In Section 5, we make this cyclicity condition explicit when the finite-dimensional irreducible representations are Kirillov-Reshetikhin modules. To conclude, we provide the structure of local Weyl modules of $Y(\mathfrak{g})$, where $\mathfrak{g}$ is simple Lie algebra of type either E or F .

2 Yangians and their representations

In this section, we recall the definition of the Yangian $Y(\mathfrak{g})$ and some results concerning its finite-dimensional representations.

Definition 2.1. Let $\mathfrak{g}$ be a simple Lie algebra over $\mathbb{C}$ with rank l and let $A = (a_{ij})_{i,j \in I}$, where $I = \{1, 2, \dots, l\}$, be its Cartan matrix. Let $D = \text{diag}(d_1, \dots, d_l)$, $d_i \in \mathbb{N}$, such that $d_1, d_2, \dots, d_l$ are co-prime and DA is symmetric. The Yangian $Y(\mathfrak{g})$ is defined to be the associative algebra with generators $x_{i,r}^\pm$, $h_{i,r}$, $i \in I$, $r \in \mathbb{Z}_{\geq 0}$, and the following defining relations:

$$\begin{aligned} [h_{i,r}, h_{j,s}] &= 0, & [h_{i,0}, x_{j,s}^\pm] &= \pm d_i a_{ij} x_{j,s}^\pm, & [x_{i,r}^+, x_{j,s}^-] &= \delta_{i,j} h_{i,r+s}, \\ [h_{i,r+1}, x_{j,s}^\pm] - [h_{i,r}, x_{j,s+1}^\pm] &= \pm \frac{1}{2} d_i a_{ij} (h_{i,r} x_{j,s}^\pm + x_{j,s}^\pm h_{i,r}), \\ [x_{i,r+1}^\pm, x_{j,s}^\pm] - [x_{i,r}^\pm, x_{j,s+1}^\pm] &= \pm \frac{1}{2} d_i a_{ij} (x_{i,r}^\pm x_{j,s}^\pm + x_{j,s}^\pm x_{i,r}^\pm), \\ \sum_{\pi} [x_{i,r_{\pi(1)}}^\pm, [x_{i,r_{\pi(2)}}^\pm, \dots, [x_{i,r_{\pi(m)}}^\pm, x_{j,s}^\pm] \cdots]] &= 0, i \neq j, \end{aligned}$$

for all sequences of non-negative integers $r_1, \dots, r_m$, where $m = 1 - a_{ij}$ and the sum is over all permutations π of $\{1, \dots, m\}$.

For the simplicity, we denote the generators of $Y(\mathfrak{sl}_2)$ by $x_k^\pm$ and h_k . In $Y(\mathfrak{g})$, for a fixed $i \in I$, $\{x_{i,r}^\pm, h_{i,r} | r \geq 0\}$ generates a subalgebra Y_i of $Y(\mathfrak{g})$ and the assignments $\frac{\sqrt{d_i}}{d_i^{k+1}} x_{i,k}^\pm \rightarrow x_k^\pm$ and $\frac{1}{d_i^{k+1}} h_{i,k} \rightarrow h_k$ define an isomorphism from Y_i to $Y(\mathfrak{sl}_2)$. We denote by Δ_{Y_i} and $\Delta_{Y(\mathfrak{g})}$ the coproducts of Y_i and $Y(\mathfrak{g})$, respectively. Let $Y^\pm$ and H be the subalgebras of $Y(\mathfrak{g})$ generated by the $x_{\alpha,k}^\pm$, for all positive roots α , and

$h_{i,k}$ for all $i \in I$ and $k \in \mathbb{Z}_{\geq 0}$, respectively. Set $N^\pm = \sum_{i,k} x_{i,k}^\pm Y^\pm$. Let $h_i(u) = 1 + h_{i,0}u^{-1} + h_{i,1}u^{-2} + \dots$ and $H_i(u) = \sum_{k=0}^{\infty} H_{i,k}u^{-k-1} := \ln(h_i(u))$.

Lemma 2.2 (Corollary 1.5, [13]).

$$[H_{i,k}, x_{j,l}^\pm] = \pm d_i a_{ij} x_{j,l+k}^\pm \pm \sum_{\substack{0 \leq s \leq k-2 \\ k+s \text{ even}}} 2^{s-k} (d_i a_{ij})^{k+1-s} \frac{\binom{k+1}{s}}{k+1} x_{j,l+s}^\pm.$$

It has been established that $Y(\mathfrak{g})$ has a Hopf algebra structure. The coproduct of $Y(\mathfrak{g})$, however, is not given explicitly in terms of the generators $x_{i,r}^\pm$ and $h_{i,r}$. Chari and Pressley proved the following proposition which is enough for our purpose.

Proposition 2.3 (Proposition 2.8, [5]).

Denote $Y(\mathfrak{g})$ by Y . Modulo $\bar{Y} \equiv (N^- Y \otimes Y N^+) \cap (Y N^- \otimes N^+ Y)$, we have

$$\begin{aligned} (i) \quad \Delta_{Y(\mathfrak{g})}(x_{i,k}^+) &\equiv x_{i,k}^+ \otimes 1 + 1 \otimes x_{i,k}^+ + \sum_{j=1}^k h_{i,j-1} \otimes x_{i,k-j}^+, \\ (ii) \quad \Delta_{Y(\mathfrak{g})}(x_{i,k}^-) &\equiv x_{i,k}^- \otimes 1 + 1 \otimes x_{i,k}^- + \sum_{j=1}^k x_{i,k-j}^- \otimes h_{i,j-1}, \\ (iii) \quad \Delta_{Y(\mathfrak{g})}(h_{i,k}) &\equiv h_{i,k} \otimes 1 + 1 \otimes h_{i,k} + \sum_{j=1}^k h_{i,j-1} \otimes h_{i,k-j}. \end{aligned}$$

We are in the position to introduce the finite-dimensional highest weight representations. In [9], Drinfeld gave the following classification of the finite-dimensional irreducible representations of $Y(\mathfrak{g})$.

Theorem 2.4.

- (i) Every irreducible finite-dimensional representation of $Y(\mathfrak{g})$ is highest weight.
- (ii) The irreducible representation $L(\mu)$ is finite-dimensional if and only if there exists an l -tuple of polynomials $\pi = (\pi_1(u), \dots, \pi_l(u))$ such that

$$\frac{\pi_i(u + d_i)}{\pi_i(u)} = 1 + \sum_{k=0}^{\infty} \mu_{i,k} u^{-k-1},$$

in the sense that the right-hand side is the Laurent expansion of the left-hand side about $u = \infty$. The polynomials $\pi_i(u)$ are called Drinfeld polynomials.

Note that every finite-dimensional highest-weight representation is also a lowest weight representation and vice-versa. The following lemma is an analogue of Lemma 4.2 in [4]. The proof has been omitted as it is similar to the proof of that Lemma.

Lemma 2.5. *Let V and W be two finite-dimensional highest weight representations of $Y(\mathfrak{g})$ with lowest and highest weight vectors v^- and w^+ , respectively. Then $v^- \otimes w^+$ generates $V \otimes W$.*

We now turn our attention to the representation theory of $Y(\mathfrak{sl}_2)$. Denote by $V_m(a)$ the finite-dimensional irreducible representation of $Y(\mathfrak{sl}_2)$ associated to the Drinfeld polynomial $(u-a)(u-(a_1+1)) \dots (u-(a+m-1))$. $V_m(a)$ is called evaluation representation of $Y(\mathfrak{sl}_2)$; as a $\mathfrak{sl}_2$ -module, $V_m(a)$ is isomorphic to the highest weight $\mathfrak{sl}_2$ -module with highest weight m .

Proposition 2.6 (Proposition 3.5,[6]). *For any $m \geq 1$, $V_m(a)$ has a basis $\{w_0, w_1, \dots, w_m\}$ on which the action of $Y(\mathfrak{sl}_2)$ is given by*

$$\begin{aligned} x_k^+ w_s &= (s+a)^k (s+1) w_{s+1}, & x_k^- w_s &= (s+a-1)^k (m-s+1) w_{s-1}, \\ h_k w_s &= ((s+a-1)^k s(m-s+1) - (s+a)^k (s+1)(m-s)) w_s. \end{aligned}$$

Call that a polynomial P is in general position with respect to another polynomial Q if for any root t of Q and any root s of P , we have $t-s \neq 1$. The following lemma is from paraphrasing Corollary 3.8 [6].

Lemma 2.7. *Let $P_1, P_2, \dots, P_m$ be polynomials, and let $V(P_i)$ be the irreducible highest weight representation whose Drinfeld polynomial is P_i . If P_j is in general position with respect to P_i ($i < j$), then $V(P_1) \otimes V(P_2) \otimes \dots \otimes V(P_m)$ is a highest weight $Y(\mathfrak{sl}_2)$ -module.*

We close this section by introducing the local Weyl module $W(\pi)$ of $Y(\mathfrak{g})$. It has been established that the category of finite-dimensional representations of $Y(\mathfrak{g})$ is not semi-simple. Therefore, it is not enough to just study its irreducible representations. In [8], the notion of local Weyl modules for classical and quantum affine algebra was introduced, and we extended this notion for Yangians [16]. The local Weyl module $W(\pi)$ has a nice universal property: any finite-dimensional highest weight representation of $Y(\mathfrak{g})$ associated to π is a quotient of $W(\pi)$.

Definition 2.8. *Let $\pi_i(u) = \prod_{j=1}^{m_i} (u - a_{i,j})$ and $\pi = (\pi_1(u), \pi_2(u), \dots, \pi_l(u))$. The local Weyl module $W(\pi)$ for $Y(\mathfrak{g})$ is defined as the module generated by a highest weight vector w_π that satisfies the following relations:*

$$x_{i,k}^+ w_\pi = 0, \quad (x_{i,0}^-)^{m_i+1} w_\pi = 0, \quad \left(h_i(u) - \frac{\pi_i(u+d_i)}{\pi_i(u)} \right) w_\pi = 0.$$

We showed in Theorem 3.8 of [16] that $\text{Dim}(W(\pi)) \leq \text{Dim}(W(\lambda_\pi))$, where $\lambda_\pi = \sum_{i=1}^l m_i \lambda_i$ and $W(\lambda_\pi)$ is the local Weyl module of the corresponding current algebra. The structure of the local Weyl modules of $Y(\mathfrak{g})$ was provided in Theorem 5.23 of [16], where $\mathfrak{g}$ is classical, and Theorem 2 [17], where $\mathfrak{g}$ is the simple Lie algebra of type G_2 . In the end of this paper, we will provide the structure of the local Weyl module of $Y(\mathfrak{g})$, when $\mathfrak{g}$ is of type E or F .

3 Braid group action

Let $\mathcal{B}$ be the braid group associated to $\mathfrak{g}$, which is the group generated by elements T_i ($i \in I$) with defining relations:

$$\begin{aligned} T_i T_j &= T_j T_i, \quad \text{if } a_{ij} = 0, \\ T_i T_j T_i &= T_j T_i T_j, \quad \text{if } a_{ij} a_{ji} = 1, \\ (T_i T_j)^2 &= (T_j T_i)^2, \quad \text{if } a_{ij} a_{ji} = 2, \\ (T_i T_j)^3 &= (T_j T_i)^3, \quad \text{if } a_{ij} a_{ji} = 3, \end{aligned}$$

where $i, j \in \{1, 2, \dots, l\}$. If $w = s_{r_1} s_{r_2} \dots s_{r_p}$ is a reduced expression of w , then denote $T_{r_1} T_{r_2} \dots T_{r_p}$ by T_w .

Recall that $\pi_{i,a}$ is the l -tuple of polynomials in u such that $\pi_i(u) = u - a$ and $\pi_j(u) = 1$. Let $\mathcal{P}$ be the set of l -tuple of rational functions in u . Then $\mathcal{P}$ is a group with generators $\pi_{i,a}$. We now define an action of a braid group of $\mathfrak{g}$ on $\mathcal{P}$.

Proposition 3.1. *let $P = (P_1(u), P_2(u), \dots, P_l(u)) \in \mathcal{P}$ and $w = s_{r_1} s_{r_2} \dots s_{r_p}$ be a reduced expression of w . The following formulas define an action of the braid group $\mathcal{B}$ of $\mathfrak{g}$ on $\mathcal{P}$: $T_w(P) = T_{r_1}(T_{r_2}(\dots T_{r_p}(P)))$ and $T_j(P)$ is defined by,*

$$\begin{aligned} (T_j(P))_i &= P_i(u), \quad \text{if } a_{ij} = 0, \\ (T_j(P))_i &= P_i(u) P_j(u - \frac{d_i}{2}), \quad \text{if } a_{ij} = -1, \\ (T_j(P))_i &= P_i(u) P_j(u - 1) P_j(u), \quad \text{if } a_{ij} = -2, \\ (T_j(P))_i &= P_i(u) P_j(u - \frac{3}{2}) P_j(u - \frac{1}{2}) P_j(u + \frac{1}{2}), \quad \text{if } a_{ij} = -3, \\ (T_j(P))_j &= \frac{1}{P_j(u - d_j)}. \end{aligned}$$

Proof. We first claim that it is enough to prove this proposition for $P = \pi_{i,a}$. Let P and Q be two l -tuple of rational functions in u . Suppose $a_{i,j} = -2$. Then $(T_j(P))_i =$

$$P_i(u)P_j(u-1)P_j(u).$$

$$\begin{aligned} (T_j(PQ))_i &= (PQ)_i(u)(PQ)_j(u-1)(PQ)_j(u) \\ &= ((P)_i(u)(Q)_i(u))((P)_j(u-1)(Q)_j(u-1))((P)_j(u)(Q)_j(u)) \\ &= ((P)_i(u)(P)_j(u-1)(P)_j(u))((Q)_i(u)(Q)_j(u-1)(Q)_j(u)) \\ &= (T_j(P)T_j(Q))_i. \end{aligned}$$

For the other cases, the proofs are similar.

Note that $T_m(\pi_{i,a}) = \pi_{i,a}$ if $i \neq m$. We next compute $T_i(\pi_{i,a})$. In what following, we may assume that $j = i + 1$ and understand $\pi_{l+1,a} = \pi_{0,a} = 1$ for any $a \in \mathbb{C}$. Without loss of generality, we use the Cartan matrix as in [3]. We claim:

- (i) If $a_{ij}a_{ji} = 1$, $T_i(\pi_{i,a}) = \pi_{i-1,a+\frac{d_i}{2}}\pi_{i,a+d_i}^{-1}\pi_{i+1,a+\frac{d_i}{2}}$ and $T_{i+1}(\pi_{i+1,a}) = \pi_{i,a+\frac{d_i}{2}}\pi_{i,a+d_i}^{-1}\pi$, where either $\pi = \pi_{i+2,a+\frac{d_{i+2}}{2}}$ if $a_{i+2,i+1} = -1$ or $\pi = \pi_{i+2,a+1}\pi_{i+2,a}$ if $a_{i+2,i+1} = -2$.
- (ii) If $a_{ij}a_{ji} = 2$,
 - (a) If $a_{ij} = -1$, then $T_i(\pi_{i,a}) = \pi_{i-1,a+\frac{d_i-1}{2}}\pi_{i,a+2}^{-1}\pi_{i+1,a+1}\pi_{i+1,a}$ and $T_j(\pi_{j,a}) = \pi_{j-1,a+\frac{d_j}{2}}\pi_{j,a+1}^{-1}\pi_{j+1,a+\frac{d_{j+1}}{2}}$.
 - (b) If $a_{ij} = -2$, then the only case is $\mathfrak{g}$ is of type C and $i = l - 1$ and $j = l$. $T_{l-1}(\pi_{l-1,a}) = \pi_{l-2,a+\frac{1}{2}}\pi_{l-1,a+1}^{-1}\pi_{l,a+1}$ and $T_l(\pi_{l,a}) = \pi_{l-1,a+1}\pi_{l-1,a}\pi_{l,a+2}^{-1}$.
- (iii) If $a_{ij}a_{ji} = 3$, then $\mathfrak{g} = G_2$, $i = 1$ and $j = 2$. $T_1(\pi_{1,a}) = \pi_{1,a+3}^{-1}\pi_{2,a+\frac{3}{2}}\pi_{2,a+\frac{1}{2}}\pi_{2,a-\frac{1}{2}}$ and $T_2(\pi_{2,a}) = \pi_{1,a+\frac{3}{2}}\pi_{2,a+1}^{-1}$.

Then it is easy to check that this action is well defined with respect to the defining relations of the Braid group; we omit the details. $\square$

4 Main results

The subalgebra generated by $x_{i,0}^\pm$ and $h_{i,0}$, $i \in I$, is isomorphic to the universal enveloping algebra $U(\mathfrak{g})$, hence $V := V(\pi)$ can be viewed as a $U(\mathfrak{g})$ -module with highest weight $\lambda_\pi = \sum_{i \in I} \text{Deg}(\pi_i)\omega_i$. The weight subspace V_{λ_π} is one-dimensional and spanned by a highest weight vector v^+ of V . Let $w = s_{r_1}s_{r_2}\dots s_{r_p}$ be a reduced expression of w in the Weyl group $\mathcal{W}$ of $\mathfrak{g}$, where s_{r_i} be a simple reflection. Suppose $s_{r_{j+1}}s_{r_{j+2}}\dots s_{r_p}(\lambda_\pi) = m_j\omega_{r_j} + \sum_{n \neq r_j} c_n\omega_n$. Then set

$$v_{w(\lambda_\pi)} = (x_{r_1,0}^-)^{m_1} (x_{r_2,0}^-)^{m_2} \dots (x_{r_p,0}^-)^{m_p} v^+.$$

In particular, when $w = w_0$, $v_{w_0(\lambda_\pi)} = v^-$, where w_0 is the longest element in $\mathcal{W}$ and v^- is a lowest weight vector of V . It is well known that the weight space $V_{w(\lambda_\pi)}$ of weight $w(\lambda_\pi)$ is one dimensional spanned by an extremal vector $v_{w(\lambda_\pi)}$ and $h_{i,k}$ acts on $v_{w(\lambda_\pi)}$ by a scalar $\mu_{ik,w}$ [5]. Denote by $\mu_{i,w}(u)$ the formal series $1 + \sum_{k \in \mathbb{Z}_{\geq 0}} \mu_{ik,w} u^{-k-1}$ and set $\Delta_w^+ = \{\alpha \in \Delta^+ : w^{-1}(\alpha) \in -\Delta^+\}$, where Δ^+ is denoted the set of positive root as the convention.

Lemma 4.1 (Remark 3.1, [5]). *There exists a polynomial $P_{i,w}(u)$ such that*

$$\mu_{i,w}(u) = \begin{cases} \frac{P_{i,w}(u+d_i)}{P_{i,w}(u)}, & \text{if } \alpha_i \notin \Delta_w^+; \\ \frac{P_{i,w}(u)}{P_{i,w}(u+d_i)}, & \text{if } \alpha_i \in \Delta_w^+. \end{cases}$$

Let V be a finite-dimensional highest weight representation of $Y(\mathfrak{sl}_2)$ whose associated polynomial is π . Let v^+ and v^- be highest and lowest weight vectors of V , respectively. It follows from Lemma 4.1 that

Corollary 4.2. $h(u)v^- = \frac{\pi(u-1)}{\pi(u)}v^-$.

Define $\sigma_j := s_{r_{j+1}}s_{r_{j+2}} \dots s_{r_p}$ and $v_{\sigma_j(\lambda_\pi)} := \left(x_{r_{j+1},0}^-\right)^{m_{j+1}} \left(x_{r_{j+2},0}^-\right)^{m_{j+2}} \dots \left(x_{r_p,0}^-\right)^{m_p} v^+$.

Lemma 4.3. $Y_{r_j}(v_{\sigma_j(\lambda_\pi)})$ is a highest weight representation of Y_{r_j} .

Proof. Suppose that $x_{r_j,0}^+ v_{\sigma_j(\lambda_\pi)} \neq 0$, then $\sigma_j(\lambda_\pi) + \alpha_{r_j}$ is a weight of V . Thus $\lambda_\pi + \sigma_j^{-1}(\alpha_{r_j})$ is a weight which does not precede λ_π , which implies that $\sigma_j^{-1}(\alpha_{r_j})$ is a negative root. However, $\ell(s_{r_j}\sigma_j) = \ell(\sigma_j) + 1$, hence $\sigma_j^{-1}(\alpha_{r_j})$ is a positive root, so we have a contradiction. Thus $x_{r_j,0}^+ v_{\sigma_j(\lambda_\pi)} = 0$. Since the weight space of weight $\sigma_j(\lambda_\pi)$ is 1-dimensional and H is a commutative subalgebra of $Y(\mathfrak{g})$, $h_{r_j,s} v_{\sigma_j(\lambda_\pi)} = c_{j,s} v_{\sigma_j(\lambda_\pi)}$, where $c_{j,s} \in \mathbb{C}$. Thus $Y_{r_j}(v_{\sigma_j(\lambda_\pi)})$ is a highest weight representation of Y_{r_j} . $\square$

The coming lemma was proved in [16] and [17].

Lemma 4.4. *Let v^+ be a highest weight representation of j -th fundamental representation $W_a(\omega_j)$ of $Y(\mathfrak{g})$. The associated polynomial $P_{i,s_j}(u)$ is listed as following:*

$$(i) \quad P_{i,s_j}(u) = 1 \text{ if } a_{ij} = 0.$$

$$(ii) \quad P_{i,s_j}(u) = u - (a + \frac{d_i}{2}) \text{ if } a_{ij} = -1.$$

$$(iii) \quad P_{i,s_j}(u) = (u - (a + 1))(u - a) \text{ if } a_{ij} = -2.$$

$$(iv) \quad P_{i,s_j}(u) = (u - (a + \frac{3}{2}))(u - (a + \frac{1}{2}))(u - (a - \frac{1}{2})) \text{ if } a_{ij} = -3.$$

Let $l(w)$ be the length of $w \in \mathcal{W}$. Suppose $l(s_i w) = l(w) + 1$. It follows from Lemma 4.3 that $h_i(u)v_{w(\lambda_\pi)} = \mu_{i,w}(u)v_{w(\lambda_\pi)}$. In the next proposition, we show that the eigenvalues $\mu_{i,w}(u)$ is determined by the braid group action on π .

Proposition 4.5. $\mu_{i,w}(u) = \frac{(T_w(\pi))_i(u+d_i)}{(T_w(\pi))_i(u)}.$

Proof. We proceed by induction on $l(w)$. If $l(w) = 0$, the claim is true since $T_w(\pi) = \pi$. Suppose that $l(w) = 1$, say $w = s_j$. Since the weight space $V_{s_j(\lambda_\pi)}$ is 1-dimensional, we may assume that $v_{s_j(\lambda_\pi)} = (x_{j,0}^-)^{m_j} v^+ \neq 0$. We are going to compute the eigenvalues $\mu_{i,k}$ of $h_{i,k}$ on the weight vector $(x_{j,0}^-)^{m_j} v^+$. Recall that $H_i(u) = \sum_{k=0}^{\infty} H_{i,k} u^{-k-1} := \ln(h_i(u))$. Denote $\sum_{r=0}^{m_j-1} (x_{j,0}^-)^r x_{j,s}^- (x_{j,0}^-)^{m_j-r-1} v^+$ by E_s and $\sum_{r=0}^{m_j-1} (x_{j,0}^-)^r [H_{i,k}, x_{j,0}^-] (x_{j,0}^-)^{m_j-r-1} v^+$ by F_k . Note that

$$\begin{aligned} H_{j,k}(x_{j,0}^-)^{m_j} v^+ &= (x_{j,0}^-)^{m_j} H_{j,k} v^+ + [H_{j,k}, (x_{j,0}^-)^{m_j}] v^+ \\ &= (x_{j,0}^-)^{m_j} H_{j,k} v^+ + \sum_{r=0}^{m_j-1} (x_{j,0}^-)^r [H_{j,k}, x_{j,0}^-] (x_{j,0}^-)^{m_j-r-1} v^+. \end{aligned}$$

Therefore, $\sum_{r=0}^{m_j-1} (x_{j,0}^-)^r [H_{j,k}, x_{j,0}^-] (x_{j,0}^-)^{m_j-r-1} v^+ = H_{j,k}(x_{j,0}^-)^{m_j} v^+ - (x_{j,0}^-)^{m_j} H_{j,k} v^+$. Denote $H_{j,k}(x_{j,0}^-)^{m_j} v^+ - (x_{j,0}^-)^{m_j} H_{j,k} v^+$ by G_k .

We show next that $F_k = \sum_{t=0}^k c_t G_t$ and each c_t does not depend on m_j . By Lemma 2.2, suppose that $[H_{i,k}, x_{j,0}^-] = \sum_{s=0}^k a_{k,s} x_{j,s}^-$ and $[H_{j,k}, x_{j,0}^-] = \sum_{s=0}^k b_{k,s} x_{j,s}^-$, where $b_{k,k} \neq 0$. Then $F_k = \sum_{s=0}^k a_{k,s} E_s$ and $G_k = \sum_{s=0}^k b_{k,s} E_s$. We note that the values of $a_{k,s}$ and $b_{k,s}$ do not depend on m_j . It is clear that $E_s = \sum_{t=0}^s b'_{s,t} G_t$. In particular, the values of $b'_{s,t}$ do not depend on m_j .

$$F_k = \sum_{s=0}^k a_{k,s} E_s = \sum_{s=0}^k a_{k,s} \left(\sum_{t=0}^s b'_{s,t} G_t \right) = \sum_{t=0}^k c_t G_t.$$

Since the coefficient c_t is only depends on both $a_{k,s}$ and $b'_{s,t}$, the coefficient c_t does not change the values with respect to m_j (the exact values of c_t are irrelevant here).

Suppose $a_{ij} = -1$. Consider the case when $V = W_a(\omega_j)$ in this paragraph. In this case, $\pi_j(u) = u - a$ and $m_j = 1$. By Lemma 4.4,

$$[h_i(u), x_{j,0}^-] v^+ = h_i(u) x_{j,0}^- v^+ = \frac{1 - (a + \frac{d_i}{2} - d_i) u^{-1}}{1 - (a + \frac{d_i}{2}) u^{-1}} x_{j,0}^- v^+.$$

Recall that $H_i(u) = \ln(h_i(u))$. Then

$$[H_i(u), x_{j,0}^-]v^+ = \left(\ln \left(1 - \left(a - \frac{d_i}{2} \right) u^{-1} \right) - \ln \left(1 - \left(a + \frac{d_i}{2} \right) u^{-1} \right) \right) x_{j,0}^- v^+.$$

Therefore, by comparing the coefficient of u^{-k-1} both sides of the above equality,

$$F_k = [H_{i,k}, x_{j,0}^-]v^+ = \frac{1}{k+1} \left(\left(a + \frac{d_i}{2} \right)^{k+1} - \left(a - \frac{d_i}{2} \right)^{k+1} \right) x_{j,0}^- v^+.$$

On the other hand, we have $h_j(u)v^+ = \frac{u-(a-d_j)}{u-a}v^+$ and $h_j(u)x_{j,0}^-v^+ = \frac{u-(a+d_j)}{u-a}x_{j,0}^-v^+$. Then

$$G_t = H_{j,t}x_{j,0}^-v^+ - x_{j,0}^-H_{j,t}v^+ = \frac{1}{t+1} \left((a-d_j)^{t+1} - (a+d_j)^{t+1} \right) x_{j,0}^- v^+.$$

Since $F_k = \sum_{t=0}^k c_t G_t$, by comparing their coefficients of $x_{j,0}^-v^+$ both sides, we obtain:

$$\frac{1}{k+1} \left(\left(a + \frac{d_i}{2} \right)^{k+1} - \left(a - \frac{d_i}{2} \right)^{k+1} \right) = \sum_{t=0}^k \frac{c_t}{t+1} \left((a-d_j)^{t+1} - (a+d_j)^{t+1} \right). \quad (4.1)$$

Now, suppose $\pi_j(u) = \prod_{s=1}^{m_j} (u - a_s)$ for $m_j \geq 2$. Thus $h_j(u)v^+ = \prod_{s=1}^{m_j} \frac{u-(a_s-d_j)}{u-a_s}v^+$ and $h_j(u)(x_{j,0}^-)^{m_j}v^+ = \prod_{s=1}^{m_j} \frac{u-(a_s+d_j)}{u-a_s}(x_{j,0}^-)^{m_j}v^+$, then

$$H_j(u)v^+ = \sum_{s=1}^{m_j} \left(\ln \left(1 - (a_s - d_j)u^{-1} \right) - \ln \left(1 - a_s u^{-1} \right) \right) v^+$$

and

$$H_j(u)(x_{j,0}^-)^{m_j}v^+ = \sum_{s=1}^{m_j} \left(\ln \left(1 - (a_s + d_j)u^{-1} \right) - \ln \left(1 - a_s u^{-1} \right) \right) (x_{j,0}^-)^{m_j}v^+.$$

Therefore,

$$G_t = H_{j,k}(x_{j,0}^-)^{m_j}v^+ - (x_{j,0}^-)^{m_j}H_{j,k}v^+ = \sum_{s=1}^{m_j} \frac{1}{t+1} \left((a_s - d_j)^{t+1} - (a_s + d_j)^{t+1} \right) (x_{j,0}^-)^{m_j}v^+.$$

$$F_k = \sum_{t=0}^k c_t G_t$$

$$\begin{aligned}
&= \sum_{t=0}^k \left(\frac{c_t}{t+1} \sum_{s=1}^{m_j} ((a_s - d_j)^{t+1} - (a_s + d_j)^{t+1}) \right) (x_{j,0}^-)^{m_j} v^+ \\
&= \sum_{s=1}^{m_j} \left(\sum_{t=0}^k \frac{c_t}{t+1} ((a_s - d_j)^{t+1} - (a_s + d_j)^{t+1}) \right) (x_{j,0}^-)^{m_j} v^+ \\
&= \sum_{s=1}^{m_j} \frac{1}{k+1} \left((a_s + \frac{d_i}{2})^{k+1} - (a_s - \frac{d_i}{2})^{k+1} \right) (x_{j,0}^-)^{m_j} v^+. \quad (\text{by (4.1)})
\end{aligned}$$

Note that the coefficient of $(x_{j,0}^-)^{m_j} v^+$ in the above computation equals the coefficient of u^{-k-1} in $\sum_{s=1}^{m_j} \ln \left(\frac{1 - (a_s - \frac{d_i}{2})u^{-1}}{1 - (a_s + \frac{d_i}{2})u^{-1}} \right)$. Thus

$$\sum_{r=0}^{m_j} (x_{j,0}^-)^r [H_i(u), x_{j,0}^-] (x_{j,0}^-)^{m_j-r-1} v^+ = \ln \left(\prod_{s=1}^{m_j} \frac{1 - (a_s - \frac{d_i}{2})u^{-1}}{1 - (a_s + \frac{d_i}{2})u^{-1}} \right) (x_{j,0}^-)^{m_j} v^+.$$

Thus we have that

$$\begin{aligned}
H_i(u)(x_{j,0}^-)^{m_j} v^+ &= \sum_{r=0}^{m_j} (x_{j,0}^-)^r [H_i(u), x_{j,0}^-] (x_{j,0}^-)^{m_j-r-1} v^+ + (x_{j,0}^-)^{m_j} H_i(u) v^+ \\
&= \ln \left(\frac{P_j(u - \frac{d_i}{2} + d_i)}{P_j(u - \frac{d_i}{2})} \right) (x_{j,0}^-)^{m_j} v^+ + \ln \left(\frac{P_i(u + d_i)}{P_i(u)} \right) (x_{j,0}^-)^{m_j} v^+ \\
&= \ln \left(\frac{P_j(u - \frac{d_i}{2} + d_i)}{P_j(u - \frac{d_i}{2})} \cdot \frac{P_i(u + d_i)}{P_i(u)} \right) (x_{j,0}^-)^{m_j} v^+.
\end{aligned}$$

Recall that $(T_j(P))_i(u) = P_i(u)P_j(u - \frac{d_i}{2})$. So

$$h_i(u)(x_{j,0}^-)^{m_j} v^+ = \frac{P_j(u - \frac{d_i}{2} + d_i)}{P_j(u - \frac{d_i}{2})} \cdot \frac{P_i(u + d_i)}{P_i(u)} (x_{j,0}^-)^{m_j} v^+ = \frac{(T_j(P))_i(u + d_i)}{(T_j(P))_i(u)} (x_{j,0}^-)^{m_j} v^+,$$

$$\text{i.e., } \mu_{i,w}(u) = \frac{(T_w(P))_i(u + d_i)}{(T_w(P))_i(u)}.$$

Similar discussions can be carried on when $a_{ij} = 0, -2$ or -3 . We omit the details.

Suppose $l(w) = n \geq 2$ and $w = s_k w'$ with $l(w') = n - 1$. Note that $T_w = T_k T_{w'}$, $T_w(\pi) = T_k(T_{w'}(\pi))$ and $v_{w(\lambda_\pi)} = (x_{k,0}^-)^{m_k} v_{w'(\lambda_\pi)}$ for some $m_k \in \mathbb{Z}_{\geq 0}$. This proof is almost the same as the above paragraphs by replacing v^+ by $v_{w'(\lambda_\pi)}$. By induction, this completes the proof of this proposition. $\square$

Remark 4.6. Suppose that $(T_{\sigma_j}(\pi))_{r_j}(u) = \prod_{r=1}^n \prod_{s_r=0}^{m_r-1} (u - (a_r + s_r))$. Since one need to normalize the generators of Y_{r_j} to satisfy the defining relations of $Y(\mathfrak{sl}_2)$, the

associated polynomial of $Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(m)})} \right)$ as a $Y(\mathfrak{sl}_2)$ -module is $\prod_{r=1}^n \prod_{s_r=0}^{m_r-1} (u - \frac{a_r + s_r}{d_i})$, which is denoted by $\widetilde{(T_w(\pi))}_{r_j}$.

Before showing the main theorem of this paper, we need the following lemma first.

Lemma 4.7 (Lemma 4.3, [5]). *Let V be an finite-dimensional irreducible representation of $Y(\mathfrak{g})$ with highest weight vector v^+ , and let $\mathfrak{t}$ be a diagram subalgebra of $\mathfrak{g}$. Then $Y(\mathfrak{t})$ -subrepresentation of V generated by v^+ is irreducible.*

From now on, we assume $s_{r_1} s_{r_2} \dots s_{r_p}$ is a reduced expression of the longest element $w_0 \in \mathcal{W}$.

Theorem 4.8. *Let $\pi^{(i)}$, $1 \leq i \leq k$, be l -tuples of polynomials in u and $V(\pi^{(i)})$ be the finite-dimensional irreducible representations of $Y(\mathfrak{g})$ whose Drinfeld polynomials are $\pi^{(i)}$. The ordered tensor product $L = V(\pi^{(1)}) \otimes V(\pi^{(2)}) \otimes \dots \otimes V(\pi^{(k)})$ is a highest weight representation if the polynomial $\widetilde{(T_{\sigma_j}(\pi^{(m)}))}_{r_j}$ is in general position with respect to $\widetilde{(\pi^{(n)})}_{r_j}$ for all $1 \leq j \leq p$ and $1 \leq m < n \leq k$.*

Proof. Let v_m^+ be the highest weight vector of $V(\pi^{(m)})$ and let v_1^- be the lowest weight vector of $V(\pi^{(1)})$. Recall that $Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(m)})} \right)$ is a highest weight representation of $Y(\mathfrak{sl}_2)$. Thus $Q := \widetilde{(T_w(\pi^{(m)}))}_{r_j}$ is a polynomial.

We prove this theorem by induction on k . For $k = 1$, $L = V(\pi^{(1)})$ is irreducible and hence it is a highest weight representation. We assume that the claim is true for all positive integers less than or equal to $k - 1$ ($k \geq 2$). By the induction hypothesis, $V(\pi^{(2)}) \otimes \dots \otimes V(\pi^{(k)})$ is a highest weight representation of $Y(\mathfrak{g})$ and its highest weight vector is $v^+ = v_2^+ \otimes \dots \otimes v_k^+$. To show that L is a highest weight representation, it suffices to show $v_1^- \otimes v^+ \in Y(\mathfrak{g}) (v_1^- \otimes v^+)$ by Lemma 2.5.

We first show that

$$Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \right) \otimes Y_{r_j} (v_2^+) \otimes \dots \otimes Y_{r_j} (v_k^+) = Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+ \right).$$

For $2 \leq n \leq k$, as $Y(\mathfrak{sl}_2)$ module, $Y_{r_j}(v_n^+)$ is isomorphic to $V(\widetilde{(\pi^{(n)})}_{r_j})$ by Lemma 4.7. Since the polynomial Q is in general position with respect to $\widetilde{(\pi^{(n)})}_{r_j}$, $W(Q) \otimes Y_{r_j} (v_2^+) \otimes \dots \otimes Y_{r_j} (v_k^+)$ is a highest weight representation of $Y(\mathfrak{sl}_2)$ by Lemma 2.7. Hence its quotient $Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \right) \otimes Y_{r_j} (v_2^+) \otimes \dots \otimes Y_{r_j} (v_k^+)$ is a highest weight module of Y_{r_j} with highest weight vector $v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+$. Thus

$$Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+ \right) \supseteq Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \right) \otimes Y_{r_j} (v_2^+) \otimes \dots \otimes Y_{r_j} (v_k^+).$$

Since $\overline{Y}$ annihilates $v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+$, by Proposition 2.3 that

$$\Delta_{Y(\mathfrak{g})}(x) \left(v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+ \right) = \Delta_{Y(\mathfrak{sl}_2)}(x) \left(v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+ \right),$$

where $x \in Y_{r_j}$. Then it is easy to see that

$$Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \otimes v_2^+ \otimes \dots \otimes v_k^+ \right) \subseteq Y_{r_j} \left(v_{\sigma_j(\lambda_{\pi(1)})} \right) \otimes Y_{r_j} (v_2^+) \otimes \dots \otimes Y_{r_j} (v_k^+).$$


Therefore the claim is true.

Since $v_{\sigma_{j-1}(\lambda_{\pi(1)})} \otimes v^+ = (x_{r_j,0}^-)^{m_j} v_{\sigma_j(\lambda_{\pi(1)})} \otimes v^+ \in Y_{r_j} (v_{\sigma_j(\lambda_{\pi(1)})}) \otimes Y_{r_j} (v^+)$, we have $v_{\sigma_{j-1}(\lambda_{\pi(1)})} \otimes v^+ \in Y_{r_j} (v_{\sigma_j(\lambda_{\pi(1)})} \otimes v^+)$. By downward induction on the subscript j of $v_{\sigma_j(\lambda_{\pi(1)})}$, $v_1^- \otimes v^+ \in Y(\mathfrak{g}) (v_1^+ \otimes v^+)$. By Lemma 2.5, $L = Y(\mathfrak{g}) (v_1^+ \otimes v^+)$ is a highest weight representation. $\square$

5 The cyclicity condition made explicit

With a specific reduced expression of w_0 , one can compute explicitly the polynomial $(T_{r_{j+1}} T_{r_{j+2}} \dots T_{r_p}(\pi^{(m)}))_{r_j}$ via the braid group action defined in Section 3. Then a cyclicity condition for the tensor product L can be obtained immediately by using Theorem 4.8. In this section, we provide a finite set of numbers at which a tensor product of Kirillov-Reshetikhin modules may fail to be cyclic. At the end of this section, we give the structure of local Weyl modules when $\mathfrak{g}$ is either of type E or of type F .

When the simple Lie algebra $\mathfrak{g}$ is classical, let the nodes of the Dynkin diagram of $\mathfrak{g}$ be numbered as in [2] and use the reduced expression of w_0 therein. When $\mathfrak{g}$ is exceptional, the nodes of the Dynkin diagram of the exceptional simple Lie algebras $\mathfrak{g}$ be numbered as following.



One reduced expression for the longest element w_0 is listed as follows.

- (a) E_6 : $u_6 = s_1 s_2 s_3 s_1 s_4 s_2 s_3 s_1 s_4 s_3 s_5 s_4 s_2 s_3 s_1 s_4 s_3 s_5 s_4 s_2 s_6 s_5 s_4 s_2 s_3 s_1 s_4 s_3 s_5 s_4 s_2 s_6 s_5 s_4 s_3 s_1$.
- (b) E_7 : $u_7 = u_6 s_7 s_6 s_5 s_4 s_2 s_3 s_1 s_4 s_3 s_5 s_4 s_2 s_6 s_5 s_4 s_3 s_1 s_7 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7$.
- (c) E_8 : $u_8 s_8 s_7 s_6 s_5 s_4 s_2 s_3 s_1 s_4 s_3 s_5 s_4 s_2 s_6 s_5 s_4 s_3 s_1 s_7 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7 s_8 s_7 s_6 s_5 s_4 s_2 s_3 s_1$
 $\times s_4 s_3 s_5 s_4 s_2 s_6 s_5 s_4 s_3 s_1 s_7 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7 s_8$.

(d) F_4 : $s_1 s_2 s_1 s_3 s_2 s_1 s_3 s_2 s_3 s_4 s_3 s_2 s_1 s_3 s_2 s_3 s_4 s_3 s_2 s_1 s_3 s_2 s_3 s_4$.

(e) G_2 : $s_2 s_1 s_2 s_1 s_2 s_1$.

Note that it is enough to consider the case when the number of tensor factors is 2. We first compute a finite set of numbers at which $V\left(\pi_{a_1,1}^{(b_1)}\right) \otimes V\left(\pi_{a_2,1}^{(b_2)}\right) = V_{a_1}(\omega_{b_1}) \otimes V_{a_2}(\omega_{b_2})$ may fail to be cyclic. Set $\{a, b..c\} = \{a, b, b+1 \dots, c-1, c\}$ and $\{a..b.c\} = \{a..b-1\} \cup \{b+1..c\}$. We omit the detailed computations for the following lemma because the computations is rudimental. In Cases 5, 6 and 7, $S(b_1, b_2) = S(b_2, b_1)$, so we omit the redundant sets.

Lemma 5.1. *Let $L = V_{a_1}(\omega_{b_1}) \otimes V_{a_2}(\omega_{b_2})$ be an ordered tensor product of fundamental representations of $Y(\mathfrak{g})$. If $a_2 - a_1 \notin S(b_1, b_2)$, then L is a highest weight representation of $Y(\mathfrak{g})$, where the set $S(b_1, b_2)$ is defined as follows:*

1. When $\mathfrak{g} = \mathfrak{sl}_{l+1}$, $S(b_1, b_2) = \left\{ \frac{|b_2 - b_1|}{2} + k \mid 1 \leq k \leq \min\{b_1, l - b_2 + 1\} \right\}$.
2. When $\mathfrak{g} = \mathfrak{so}(2l + 1, \mathbb{C})$,
 - (a) $S(b_1, b_2) = \{|b_1 - b_2| + 2 + 2r, 2l - (b_1 + b_2) + 1 + 2r\}$, where $1 \leq b_1, b_2 \leq l - 1$ and $0 \leq r < \min\{b_1, b_2\}$;
 - (b) $S(l, b_2) = \{l - b_2 + 2 + 2r \mid 0 \leq r < b_2, 1 \leq b_2 \leq l - 1\}$;
 - (c) $S(b_1, l) = \{l - b_1 + 1 + r, l - b_1 + r \mid 0 \leq r < b_1, 1 \leq b_1 \leq l - 1\}$;
 - (d) $S(l, l) = \{1, 3, \dots, 2l - 1\}$.
3. When $\mathfrak{g} = \mathfrak{sp}(2l, \mathbb{C})$,
 - (a) $S(b_1, b_2) = \left\{ \frac{|b_1 - b_2|}{2} + 1 + r, l + 2 + r - \frac{b_1 + b_2}{2} \mid 0 \leq r < \min\{b_1, b_2\} \right\}$, where $1 \leq b_1, b_2 \leq l - 1$;
 - (b) $S(l, b_2) = \left\{ \frac{l - b_2 + 1}{2} + 1 + r, \frac{l - b_2 - 1}{2} + 1 + r \mid 0 \leq r < b_2, 1 \leq b_2 \leq l - 1 \right\}$;
 - (c) $S(b_1, l) = \left\{ \frac{l - b_1 + 1}{2} + 2 + r \mid 0 \leq r < b_1, 1 \leq b_1 \leq l - 1 \right\}$;
 - (d) $S(l, l) = \{2, 3, \dots, l + 1\}$.
4. When $\mathfrak{g} = \mathfrak{so}(2l, \mathbb{C})$,
 - (a) $S(b_1, b_2) = \left\{ \frac{|b_1 - b_2|}{2} + 1 + r, l + r - \frac{b_1 + b_2}{2} \right\}$, where $1 \leq b_1, b_2 \leq l - 2$ and $0 \leq r < \min\{b_1, b_2\}$;
 - (b) $S(l - 1, b_2) = S(l, b_2) = S(b_2, l - 1) = S(b_2, l) = \left\{ \frac{l - 1 - b_2}{2} + 1 + r \mid 0 \leq r < b_2, 1 \leq b_2 \leq l - 2 \right\}$;

(c) $S(l-1, l) = S(l, l-1) = \{2, 4, \dots, l-2+\bar{l}\}$, where $\bar{l} = 0$ if l is even and let $\bar{l} = 1$ if l is odd;

(d) $S(l-1, l-1) = S(l, l) = \{1, 3, \dots, l-1-\bar{l}\}$.

5. When $\mathfrak{g} = E_6$,

$$\begin{array}{lll}
S(1, 1) = \{1, 4\} & S(1, 2) = \{\frac{5}{2}, \frac{9}{2}\} & S(1, 3) = \{\frac{3}{2}, \frac{7}{2}, \frac{9}{2}\} \\
S(1, 4) = \{2, 3, 4, 5\} & S(1, 5) = \{\frac{5}{2}, \frac{7}{2}, \frac{11}{2}\} & S(1, 6) = \{3, 6\} \\
S(2, 2) = \{1, 3, 4, 6\} & S(2, 3) = \{2, 3, 4, 5\} & S(2, 4) = \{\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}\} \\
S(2, 5) = \{2, 3, 4, 5\} & S(2, 6) = \{\frac{5}{2}, \frac{9}{2}\} & S(3, 3) = \{1, 2, 3, 4, 5\} \\
S(3, 4) = \{\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}\} & S(3, 5) = \{2, 3, 4, 5, 6\} & S(3, 6) = \{\frac{5}{2}, \frac{7}{2}, \frac{11}{2}\} \\
S(4, 4) = \{1, 2, 3, 4, 5, 6\} & S(4, 5) = \{\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}\} & S(4, 6) = \{2, 3, 4, 5\} \\
S(5, 5) = \{1, 2, 3, 4, 5\} & S(5, 6) = \{\frac{3}{2}, \frac{7}{2}, \frac{9}{2}\} & S(6, 6) = \{1, 4\}.
\end{array}$$

6. When $\mathfrak{g} = E_7$,

$$\begin{array}{lll}
S(1, 1) = \{1, 4, 6, 9\}, & S(1, 2) = \{\frac{5}{2}, \frac{9}{2}, \frac{11}{2}, \frac{15}{2}\}, & S(1, 3) = \{\frac{3}{2}, \frac{7}{2}, \frac{13}{2}, \frac{17}{2}\}, \\
S(1, 4) = \{2..8\}, & S(1, 5) = \{\frac{5}{2}.. \frac{15}{2}\}, & S(1, 6) = \{3, 4, 6, 7\}, \\
S(1, 7) = \{\frac{7}{2}, \frac{13}{2}\}, & S(2, 2) = \{1, 3..7, 9\}, & S(2, 3) = \{2..8\}, \\
S(2, 4) = \{\frac{3}{2}.. \frac{17}{2}\}, & S(2, 5) = \{2..8\}, & S(2, 6) = \{\frac{5}{2}.. \frac{15}{2}\}, \\
S(2, 7) = \{3, 5, 7\}, & S(3, 3) = \{1..9\}, & S(3, 4) = \{\frac{3}{2}.. \frac{17}{2}\}, \\
S(3, 5) = \{2..8\}, & S(3, 6) = \{\frac{5}{2}.. \frac{15}{2}\}, & S(3, 7) = \{3, 4, 6, 7\}, \\
S(4, 4) = \{1..9\}, & S(4, 5) = \{\frac{3}{2}, .. \frac{17}{2}\}, & S(4, 6) = \{2..8\}, \\
S(4, 7) = \{\frac{5}{2}.. \frac{15}{2}\}, & S(5, 5) = \{1..9\}, & S(5, 6) = \{\frac{3}{2}.. \frac{17}{2}\}, \\
S(5, 7) = \{2, 4, 5, 6, 8\}, & S(6, 6) = \{1, 2, 4, 5, 6, 8, 9\}, & S(6, 7) = \{\frac{3}{2}, \frac{9}{2}, \frac{11}{2}, \frac{17}{2}\}, \\
S(7, 7) = \{1, 5, 9\}.
\end{array}$$

7. When $\mathfrak{g} = E_8$,

$$\begin{aligned}
S(1, 1) &= \{1, 4, 6, 7, 9, 10, 12, 15\}, & S(1, 2) &= \{\frac{5}{2} \dots \frac{\hat{7}}{2} \dots \frac{\hat{25}}{2} \dots \frac{27}{2}\}, \\
S(1, 3) &= \{\frac{3}{2} \dots \frac{\hat{27}}{2} \dots \frac{29}{2}\}, & S(1, 4) &= \{2..14\}, \\
S(1, 5) &= \{\frac{5}{2} \dots \frac{27}{2}\}, & S(1, 6) &= \{3..13\}, \\
S(1, 7) &= \{\frac{7}{2} \dots \frac{\hat{11}}{2} \dots \frac{\hat{21}}{2} \dots \frac{25}{2}\}, & S(1, 8) &= \{4, 7, 9, 12\}, \\
S(2, 2) &= \{1..2..14..15\}, & S(2, 3) &= \{2..14\}, \\
S(2, 4) &= \{\frac{3}{2} \dots \frac{29}{2}\}, & S(2, 5) &= \{2..14\}, \\
S(2, 6) &= \{\frac{5}{2} \dots \frac{27}{2}\}, & S(2, 7) &= \{3..13\}, \\
S(2, 8) &= \{\frac{7}{2}, \frac{11}{2}, \frac{15}{2}, \frac{17}{2}, \frac{21}{2}, \frac{25}{2}\}, & S(3, 3) &= \{1..15\}, \\
S(3, 4) &= \{\frac{3}{2} \dots \frac{29}{2}\}, & S(3, 5) &= \{2..14\}, \\
S(3, 6) &= \{\frac{5}{2} \dots \frac{27}{2}\}, & S(3, 7) &= \{3..13\}, \\
S(3, 8) &= \{\frac{7}{2} \dots \frac{\hat{11}}{2} \dots \frac{\hat{21}}{2} \dots \frac{25}{2}\}, & S(4, 4) &= \{1..15\}, \\
S(4, 5) &= \{\frac{3}{2} \dots \frac{29}{2}\}, & S(4, 6) &= \{2..14\}, \\
S(4, 7) &= \{\frac{5}{2} \dots \frac{27}{2}\}, & S(4, 8) &= \{3..13\}, \\
S(5, 5) &= \{1..15\}, & S(5, 6) &= \{\frac{3}{2} \dots \frac{29}{2}\}, \\
S(5, 7) &= \{2..14\}, & S(5, 8) &= \{\frac{5}{2} \dots \frac{\hat{7}}{2} \dots \frac{\hat{25}}{2} \dots \frac{27}{2}\}, \\
S(6, 6) &= \{1..15\}, & S(6, 7) &= \{\frac{3}{2}, \frac{7}{2} \dots \frac{15}{2} \dots \frac{25}{2} \dots \frac{29}{2}\}, \\
S(6, 8) &= \{2, 5..8..11, 14\}, & S(7, 7) &= \{1, 2, 5..8..11, 14, 15\}, \\
S(7, 8) &= \{\frac{3}{2}, \frac{11}{2}, \frac{13}{2}, \frac{19}{2}, \frac{21}{2}, \frac{29}{2}\}, & S(8, 8) &= \{1, 6, 10, 15\}.
\end{aligned}$$

8. When $\mathfrak{g} = F_4$,

$$\begin{aligned}
S(1, 1) &= \{1, 4, 5, 8\}, & S(1, 2) &= \{2..7\}, & S(1, 3) &= \{2..7\}, \\
S(1, 4) &= \{\frac{5}{2}, \frac{7}{2}, \frac{11}{2}, \frac{13}{2}\}, & S(2, 1) &= S(1, 2), & S(2, 2) &= \{1..8\}, \\
S(2, 3) &= \{1..8\}, & S(2, 4) &= \{\frac{3}{2} \dots \frac{15}{2}\}, & S(3, 1) &= \{3, 4, 6, 7\}, \\
S(3, 2) &= \{2..\hat{4}..8\}, & S(3, 3) &= \{1..9\}, & S(3, 4) &= \{\frac{3}{2} \dots \frac{\hat{5}}{2} \dots \frac{\hat{15}}{2} \dots \frac{17}{2}\}, \\
S(4, 1) &= \{\frac{7}{2}, \frac{13}{2}\}, & S(4, 2) &= \{\frac{5}{2}, \frac{9}{2}, \frac{11}{2}, \frac{15}{2}\}, & S(4, 3) &= S(3, 4), \\
S(4, 4) &= \{1, 4, 6, 9\}.
\end{aligned}$$

9. When $\mathfrak{g} = G_2$,

$$\begin{aligned}
S(1, 1) &= \{3, 4, 5, 6\}, & S(1, 2) &= \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}\}, \\
S(2, 1) &= \{\frac{9}{2}, \frac{13}{2}\}, & S(2, 2) &= \{1, 3, 4, 6\}.
\end{aligned}$$

Let $S = \{a_1, \dots, a_n\}$. Define $S + r = \{a_1 + r, \dots, a_n + r\}$.

Proposition 5.2. *An ordered tensor product $L = V\left(\pi_{a_1, m_1}^{(b_1)}\right) \otimes V\left(\pi_{a_2, m_2}^{(b_2)}\right)$ is a highest weight representation if $a_2 - a_1 \notin \bigcup_{s=0}^{m_2-1} \left(\bigcup_{r=0}^{m_1-1} (S(b_1, b_2) - s + r) \right)$, where $S(b_1, b_2)$ is defined as in Lemma 5.1.*

Proof. It follows from Proposition 3.1 that $T_w(\pi_{a_1, m_1}^{(b_1)}) = \prod_{r=0}^{m_1-1} T_w(\pi_{a_1+r, 1}^{(b_1)})$. It is easy to see that $\left(\widetilde{T_w(\pi_{a_1, m_1}^{(b_1)})}\right)_{b_2}$ is in general position with respect to $\left(\widetilde{\pi_{a_2, m_2}^{(b_2)}}\right)_{b_2}$ if and only if for each $0 \leq r \leq m_1 - 1$ and $0 \leq s \leq m_2 - 1$, $\left(\widetilde{T_w(\pi_{a_1+r, 1}^{(b_1)})}\right)_{b_2}$ is in general position with respect to $\left(\widetilde{\pi_{a_2+s, 1}^{(b_2)}}\right)_{b_2}$. By the Lemma 5.1, $(a_2 + s) - (a_1 + r) \notin S(b_1, b_2)$, which is equivalent to $a_2 - a_1 \notin \bigcup_{s=0}^{m_2-1} \left(\bigcup_{r=0}^{m_1-1} (S(b_1, b_2) - s + r) \right)$. $\square$

Note that the dimension of the local Weyl module $W(\omega_i)$ for the current algebra $\mathfrak{g}[t]$ and that of local Weyl module $W(\pi_{i,a})$ for $Y(\mathfrak{g})$ are the same. It follows from Proposition 5.2 and Theorem 3.7 in [16] that

Theorem 5.3. *Let $\mathfrak{g}$ be a complex simple Lie algebra. Let $\pi = (\pi_1(u), \dots, \pi_l(u))$, where $\pi_i(u) = \prod_{j=1}^{m_i} (u - a_{i,j})$. Let S be the multiset of the roots of these polynomials. Let $a_1 = a_{i,j}$ be one of the numbers in S with the maximal real part and let $b_1 = i$. Inductively, let $a_r = a_{s,t}$ ($r \geq 2$) be one of the numbers in $S \setminus \{a_1, \dots, a_{r-1}\}$ ($r \geq 2$) with the maximal real part and $b_r = s$. Let $k = m_1 + \dots + m_l$. Then the ordered tensor product $L = V_{a_1}(\omega_{b_1}) \otimes V_{a_2}(\omega_{b_2}) \otimes \dots \otimes V_{a_k}(\omega_{b_k})$ is a highest weight representation of $Y(\mathfrak{g})$, and its associated polynomial is π . Moreover, the local Weyl module $W(\pi)$ is isomorphic to L .*

References

- [1] T. Akasaka and M. Kashiwara, “Finite-dimensional representations of quantum affine algebras.” Publ. Res. Inst. Math. Sci. 33 (1997): 839–867.
- [2] G. Benkart, S. Kang, S. Oh and E. Park, “Construction of Irreducible Representations over Khovanov-Lauda-Rouquier Algebras of Finite Classical Type.” arXiv:1108.1048v3.
- [3] R. Carter, *Lie Algebras of Finite and Affine Type*, Cambridge Studies in Advanced Mathematics, 2005.
- [4] V. Chari, “Braid group actions and tensor products.” Int. Math. Res. Not. (2002), no. 7: 357–382.
- [5] V. Chari and A. Pressley, “Fundamental representations of Yangians and singularities of R -matrices.” J. Reine angew. Math. 417 (1991): 87–128.

- [6] V. Chari and A. Pressley, “*Yangians: their Representation and Characters.*” *Acta Applicandae Mathematica*, 44 (1996): 39-58.
- [7] V. Chari and A. Pressley, “*Yangians, integrable quantum systems and Dorey’s rule.*” *Commun. Math. Phys.* 181 (1996): 265-302.
- [8] V. Chari and A. Pressley, “*Weyl modules for classical and quantum affine algebras.*” *Represent. Theory* 5 (2001): 191–223.
- [9] V. Drinfeld, “*A new realization of Yangians and of quantum affine algebras (Russian).*” *Dokl. Akad. Nauk SSSR*. 296 (1987), no. 1: 13–17; translation in *Soviet Math. Dokl.* **36** (1988): 212–216.
- [10] E. Frenkel and E. Mukhin, “*Combinatorics of q -characters of finite-dimensional representations of quantum affine algebras.*” *Comm. Math. Phys.* 216 (2001): 23–57.
- [11] S. Gautam and L. Valerio Toledano, “*Meromorphic Kazhdan-Lusztig equivalence for Yangians and quantum loop algebras.*” arXiv: 1403.5251, 2014.
- [12] M. Kashiwara, “*On level zero representations of quantized affine algebras.*” *Duke Math. J.* 112 (2002): 117–195.
- [13] S. Z. Levendorskii, “*On generators and defining relations of Yangians.*” *Journal of Geometry and Physics* 12 (1992): 1–11.
- [14] A. Molev, V. Tolstony and R. Zhang, “*On irreducibility of tensor products of evaluation modules for the quantum affine algebras.*” *J. Phys. A.* 37 (2003): 2385–2399.
- [15] H. Nakajima, “*Quiver varieties and finite-dimensional representations of quantum affine algebras.*” *J. Amer. Math. Soc.* 14 (2001): 145–238.
- [16] Y. Tan and N. Guay, “*Local Weyl modules and cyclicity of tensor products for Yangians.*” *J. Algebra.* 432 (2015), 228–251.
- [17] Y. Tan, “*Local Weyl modules and cyclicity of the tensor products for Yangian of type G_2 .*” DOI:10.1007/s10468-015-9532-0.
- [18] M. Varagnolo, “*Quiver varieties and Yangians.*” *Lett. Math. Phys.* 53 (2000): 273–283.
- [19] M. Varagnolo and E. Vasserot, “*Standard modules of quantum affine algebras.*” *Duke Math. J.* 111 (2002): 509–533.